

TITLE OF THE INVENTION

Nucleotide sequences which code for the metR and metZ genes

BACKGROUND OF THE INVENTION

5 Field of the Invention

The invention provides nucleotide sequences from coryneform bacteria which code for the metR and metZ genes and a process for the fermentative preparation of amino acids, in particular L-methionine, by attenuation of the metR and/or metZ gene.

10 Description of the Related Art

L-Amino acids, in particular methionine, are used in human medicine and in the pharmaceuticals industry, in the foodstuffs industry and very particularly in animal nutrition.

It is known that amino acids are prepared by fermentation from strains of coryneform bacteria, in particular *Corynebacterium glutamicum*. Because of their great importance, work is constantly being undertaken to improve the preparation process. Improvements to the process can relate to fermentation measures, such as stirring and supply of oxygen, or to the composition of the nutrient media, such as, the sugar concentration during the fermentation, or to the working up of the product by, for example, ion exchange chromatography, or to the intrinsic output properties of the microorganism itself.

25 Methods of mutagenesis, selection and mutant selection are used to improve the output properties of these microorganisms. Strains which are resistant to antimetabolites or are auxotrophic for metabolites of regulatory importance and which produce amino acids, such as e.g. L-methionine, are obtained  
30 in this manner.

Recombinant DNA techniques have also been employed for some years for improving of Corynebacterium strains which produce L-amino acids, by amplifying individual amino acid biosynthesis genes and investigating their effect on the amino acid production.

#### SUMMARY OF THE INVENTION

One object of the present invention is to provide new measures for improved fermentative preparation of amino acids, in particular L-methionine.

Where L-amino acids or amino acids are mentioned in the following, this means one or more amino acids, including their salts, chosen from the group consisting of L-asparagine, L-threonine, L-serine, L-glutamate, L-glycine, L-alanine, L-cysteine, L-valine, L-methionine, L-isoleucine, L-leucine, L-tyrosine, L-phenylalanine, L-histidine, L-lysine, L-tryptophan and L-arginine.

When L-methionine or methionine are mentioned in the following, the salts, such as methionine hydrochloride or methionine sulfate are also meant.

The invention provides isolated polynucleotides from coryneform bacteria, which comprise the polynucleotide sequences which code for the metR and/or metZ genes, chosen from the group consisting of

- a) polynucleotide which is at least 70% identical to a polynucleotide which codes for a polypeptide which comprises the amino acid sequence of SEQ ID No. 2,
- b) polynucleotide which is at least 70% identical to a polynucleotide which codes for a polypeptide which comprises the amino acid sequence of SEQ ID No. 3,
- c) polynucleotide which codes for a polypeptide which comprises an amino acid sequence which is at least 70% identical to the amino acid sequence of SEQ ID No. 2,

d) polynucleotide which codes for a polypeptide which comprises an amino acid sequence which is at least 70% identical to the amino acid sequence of SEQ ID No. 3,

5 e) polynucleotide which is complementary to the polynucleotides of a), b) c) or d), and

f) polynucleotide comprising at least 15 successive nucleotides of the polynucleotide sequence of a), b), c), d) or e),

10 and the corresponding polypeptides according to a) or c) having the enzymatic activity of the transcription activator MetR and the polypeptides according to b) or d) having the enzymatic activity of O-succinylhomoserine sulphydrylase (MetZ).

15 The invention also provides the above-mentioned polynucleotides, as DNAs which are capable of replication, comprising:

- 20 (i) the nucleotide sequence shown in SEQ ID No. 1, or
- (ii) at least one sequence which corresponds to sequence (i) within the range of the degeneration of the genetic code, or
- (iii) at least one sequence which hybridizes with the sequences complementary to sequences (i) or (ii), and optionally
- (iv) sense mutations of neutral function in (i).

25 The invention also provides:

a DNA which is capable of replication and comprises the nucleotide sequence as shown in SEQ ID No. 1,

30 a polynucleotide that codes for a polypeptide which comprises the amino acid sequence as shown in SEQ ID No. 2 or SEQ ID No. 3,

a vector containing parts of the polynucleotide according to the invention, but at least 15 successive nucleotides of the sequence claimed

and coryneform bacteria in which the metR gene and/or the metZ gene is or are attenuated, in particular by deletion, insertion or base exchange.

The invention also provides polynucleotides which are obtained by screening a corresponding gene library of a coryneform bacterium, which comprises the complete gene or parts thereof by means of hybridization, with a probe which comprises the sequence of the polynucleotide according to the invention according to SEQ ID No. 1 or a fragment thereof, and isolation of the polynucleotide sequence mentioned.

#### BRIEF DESCRIPTION OF THE FIGURES

Fig. 1 shows plasmid pK18mobsacBdeltametRmetZ.

#### DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

Polynucleotides according to the invention are suitable as hybridization probes for RNA, cDNA and DNA, in order to isolate, in the full length, nucleic acids, or polynucleotides or genes which code for the transcription activator MetR and/or O-succinylhomoserine sulfhydrylase or to isolate those nucleic acids or polynucleotides or genes which have a high similarity or homology with the sequence of the transcription activator MetR gene and/or that of the O-succinylhomoserine sulfhydrylase gene.

Polynucleotides according to the invention are furthermore suitable as primers with the aid of which DNA of genes that code for the transcription activator MetR and/or O-succinylhomoserine sulfhydrylase can be prepared by the polymerase chain reaction (PCR).

Such oligonucleotides that serve as probes or primers comprise at least 30, preferably at least 20, very particularly at least 15 successive nucleotides. Oligonucleotides which have

a length of at least 40 or 50 nucleotides are also suitable. Oligonucleotides with a length of at least 100, 150, 200, 250 or 300 nucleotides are optionally also suitable.

"Isolated" means separated out of its natural environment.

5 "Polynucleotide" in general relates to polyribonucleotides and polydeoxyribonucleotides, it being possible for these to be non-modified RNA or DNA or modified RNA or DNA.

10 The polynucleotides according to the invention include a polynucleotide according to SEQ ID No. 1 or a fragment prepared therefrom and also those which are at least 70%, preferably at least 80% and in particular at least 90% to 95% identical to the polynucleotide according to SEQ ID No. 1 or a fragment prepared therefrom.

15 The polypeptides according to the invention include the polypeptides according to SEQ ID No. 2 and SEQ ID No. 3, in particular those with the biological activity of the transcription activator MetR and of O-succinylhomoserine sulfhydrylase, and also those which are at least 70%, preferably at least 80%, and in particular which are at least 90% to 95% identical to the polypeptides according to SEQ ID No. 2 and SEQ ID No. 3 and have the activities mentioned.

"Polypeptides" are understood as meaning peptides or proteins which comprise two or more amino acids bonded via peptide bonds.

25 The invention moreover provides a process for the fermentative preparation of amino acids, in particular methionine, using coryneform bacteria which in particular already produce the amino acids, and in which the nucleotide sequences which code for the metR gene and/or for the metZ gene are attenuated, in particular eliminated or expressed at a low level.

30 The term "attenuation" in this connection describes the reduction or elimination of the intracellular activity of one or more enzymes (proteins) in a microorganism which are coded by the corresponding DNA, for example by using a weak promoter

or using a gene or allele which codes for a corresponding enzyme with a low activity or inactivates the corresponding gene or enzyme (protein), and optionally combining these measures.

- 5 By attenuation measures, the activity or concentration of the corresponding protein is in general reduced to 0 to 50%, 0 to 25%, 0 to 10% or 0 to 5% of the activity or concentration of the wild-type protein.

10 The microorganisms which the present invention provides can prepare L-amino acids, in particular methionine, from glucose, sucrose, lactose, fructose, maltose, molasses, starch, cellulose or from glycerol and ethanol. They can be representatives of coryneform bacteria, in particular of the genus *Corynebacterium*. Of the genus *Corynebacterium*, there may be mentioned in particular the species *Corynebacterium glutamicum*, which is known among experts for its ability to produce L-amino acids.

Suitable strains of the genus *Corynebacterium*, in particular of the species *Corynebacterium glutamicum* (*C. glutamicum*), are in particular the known wild-type strains

*Corynebacterium glutamicum* ATCC13032  
*Corynebacterium acetoglutamicum* ATCC15806  
*Corynebacterium acetoacidophilum* ATCC13870  
*Corynebacterium melassecola* ATCC17965  
25 *Corynebacterium thermoaminogenes* FERM BP-1539  
*Brevibacterium flavum* ATCC14067  
*Brevibacterium lactofermentum* ATCC13869 and  
*Brevibacterium divaricatum* ATCC14020

or L-amino acid-producing mutants or strains prepared  
30 therefrom, such as, for example, the L-methionine-producing strain

*Corynebacterium glutamicum* ATCC21608.

The new metR and metZ genes from *C. glutamicum* which code for the transcription activator MetR and the enzyme O-succinylhomoserine sulfhydrylase have been isolated.

To isolate the metR gene, the metZ gene or also other genes of *C. glutamicum*, a gene library of this microorganism is first set up in *Escherichia coli* (*E. coli*). The setting up of gene libraries is described in generally known textbooks and handbooks. The textbook by Winnacker: *Gene und Klone, Eine Einführung in die Gentechnologie* (Verlag Chemie, Weinheim, Germany, 1990), or the handbook by Sambrook et al.: *Molecular Cloning, A Laboratory Manual* (Cold Spring Harbor Laboratory Press, 1989) may be mentioned as an example. A well-known gene library is that of the *E. coli* K-12 strain W3110 set up in  $\lambda$  vectors by Kohara et al. (*Cell* 50, 495-508 (1987)). Bathe et al. (*Molecular and General Genetics*, 252:255-265, 1996) describe a gene library of *C. glutamicum* ATCC13032, which was set up with the aid of the cosmid vector SuperCos I (Wahl et al., 1987, *Proceedings of the National Academy of Sciences USA*, 84:2160-2164) in the *E. coli* K-12 strain NM554 (Raleigh et al., 1988, *Nucleic Acids Research* 16:1563-1575). Börmann et al. (*Molecular Microbiology* 6(3), 317-326 (1992)) in turn describe a gene library of *C. glutamicum* ATCC13032 using the cosmid pH79 (Hohn and Collins, *Gene* 11, 291-298 (1980)). To prepare a gene library of *C. glutamicum* in *E. coli* it is also possible to use plasmids such as pBR322 (Bolivar, *Life Sciences*, 25, 807-818 (1979)) or pUC9 (Vieira et al., 1982, *Gene*, 19:259-268). Suitable hosts are, in particular, those *E. coli* strains which are restriction- and recombination-defective. An example of these is the strain DH5 $\alpha$ mc<sup>r</sup>, which has been described by Grant et al. (*Proceedings of the National Academy of Sciences USA*, 87 (1990) 4645-4649). The long DNA fragments cloned with the aid of cosmids can in turn be subcloned in the usual vectors suitable for sequencing and then sequenced, as is described e.g. by Sanger et al. (*Proceedings of the National Academy of Sciences of the United States of America*, 74:5463-5467, 1977).

5 The resulting DNA sequences can then be investigated with  
known algorithms or sequence analysis programs, such as that  
of Staden (Nucleic Acids Research 14, 217-232(1986)), that of  
Marck (Nucleic Acids Research 16, 1829-1836 (1988)) or the GCG  
program of Butler (Methods of Biochemical Analysis 39, 74-97  
(1998)).

10 The new DNA sequence of *C. glutamicum* which codes for the metR  
and metZ genes and which, as SEQ ID No. 1, is a constituent of  
the present invention has been found. The amino acid sequence  
of the corresponding proteins has furthermore been derived  
from the present DNA sequence by the methods described above.  
The resulting amino acid sequences of the metR and metZ gene  
products are shown in SEQ ID No. 2 and 3.

15 Coding DNA sequences which result from SEQ ID No. 1 by the  
degeneracy of the genetic code are also a constituent of the  
invention. Conservative amino acid exchanges, such as e.g.  
exchange of glycine for alanine or of aspartic acid for  
glutamic acid in proteins, are furthermore known among experts  
as "sense mutations" which do not lead to a fundamental change  
in the activity of the protein, i.e. they are of neutral  
function.

20 It is furthermore known that changes at the N and/or C  
terminus of a protein must not substantially impair and may  
even stabilize the function thereof. Information in this  
context can be found in Ben-Bassat et al. (Journal of  
25 Bacteriology 169:751-757 (1987)), in O'Regan et al. (Gene  
77:237-251 (1989)), in Sahin-Toth et al. (Protein Sciences  
3:240-247 (1994)), in Hochuli et al. (Bio/Technology 6:1321-  
1325 (1988)) and in known textbooks of genetics and molecular  
30 biology. Amino acid sequences which result in a corresponding  
manner from SEQ ID No. 2 or SEQ ID No. 3 are also a  
constituent of the invention.

35 In the same way, DNA sequences which hybridize with SEQ ID No.  
1 or parts of SEQ ID No. 1 are a constituent of the invention.  
Finally, DNA sequences which are prepared by the polymerase  
chain reaction (PCR) using primers which result from SEQ ID



No. 1 are a constituent of the invention. Such oligonucleotides typically have a length of at least 15 nucleotides.

Instructions for identifying DNA sequences by means of hybridization can be found in the handbook "The DIG System Users Guide for Filter Hybridization" from Boehringer Mannheim GmbH (Mannheim, Germany, 1993) and in Liebl et al. (International Journal of Systematic Bacteriology (1991) 41: 255-260). The hybridization takes place under stringent conditions, i.e. only hybrids in which the probe and target sequence, i.e. the polynucleotides treated with the probe, are at least 70% identical are formed. It is known that the stringency of the hybridization, including the washing steps, is influenced or determined by varying the buffer composition, the temperature and the salt concentration. The hybridization reaction is preferably carried out under a relatively low stringency compared with the washing steps (Hybaid Hybridisation Guide, Hybaid Limited, Teddington, UK, 1996).

A 5x SSC buffer at a temperature of approx. 50 - 68°C, for example, can be employed for the hybridization reaction. Probes can also hybridize here with polynucleotides which are less than 70% identical to the sequence of the probe. Such hybrids are less stable and are removed by washing under stringent conditions. This can be achieved, for example, by lowering the salt concentration to 2x SSC and optionally subsequently 0.5x SSC (The DIG System User's Guide for Filter Hybridisation, Boehringer Mannheim, Mannheim, Germany, 1995) a temperature of approx. 50 - 68°C being established. It is optionally possible to lower the salt concentration to 0.1x SSC. Polynucleotide fragments which are, for example, at least 70% or at least 80% or at least 90% to 95% identical to the sequence of the probe employed can be isolated by increasing the hybridization temperature stepwise from 50 to 68°C in steps of approx. 1 - 2°C. Further instructions on hybridization are obtainable on the market in the form of so-called kits (e.g. DIG Easy Hyb from Roche Diagnostics GmbH, Mannheim, Germany, Catalogue No. 1603558).

Instructions for amplification of DNA sequences with the aid of the polymerase chain reaction (PCR) can be found in the handbook by Gait: Oligonucleotide Synthesis: A Practical Approach (IRL Press, Oxford, UK, 1984) and in Newton and Graham: PCR (Spektrum Akademischer Verlag, Heidelberg, Germany, 1994).

It has been found that coryneform bacteria produce amino acids, in particular L-methionine, in an improved manner after attenuation of the metR and/or metZ gene.

- 10 To achieve an attenuation, either the expression of the metR and/or of the metZ gene or the catalytic properties of the enzyme proteins can be reduced or eliminated. The two measures can optionally be combined.

15 The reduction in gene expression can take place by suitable culturing or by genetic modification (mutation) of the signal structures of gene expression. Signal structures of gene expression are, for example, repressor genes, activator genes, operators, promoters, attenuators, ribosome binding sites, the start codon and terminators. The expert can find information on this e.g. in the patent application WO 96/15246, in Boyd and Murphy (Journal of Bacteriology 170: 5949 (1988)), in Voskuil and Chambliss (Nucleic Acids Research 26: 3548 (1998)), in Jensen and Hammer (Biotechnology and Bioengineering 58: 191 (1998)), in Pátek et al. (Microbiology 142: 1297 (1996)),  
20 Vasicova et al. (Journal of Bacteriology 181: 6188 (1999)) and in known textbooks of genetics and molecular biology, such as e.g. the textbook by Knippers ("Molekulare Genetik", 6th edition, Georg Thieme Verlag, Stuttgart, Germany, 1995) or that by Winnacker ("Gene und Klone", VCH Verlagsgesellschaft,  
25 Weinheim, Germany, 1990).

Mutations which lead to a change or reduction in the catalytic properties of enzyme proteins are known from the prior art; examples which may be mentioned are the works by Qiu and Goodman (Journal of Biological Chemistry 272: 8611-8617  
35 (1997)), Sugimoto et al. (Bioscience Biotechnology and Biochemistry 61: 1760-1762 (1997)) and Möckel ("Die

Threonindehydratase aus Corynebacterium glutamicum: Aufhebung der allosterischen Regulation und Struktur des Enzyms", Reports from the Jülich Research Center, Jül-2906, ISSN09442952, Jülich, Germany, 1994). Summarizing

5. descriptions can be found in known textbooks of genetics and molecular biology, such as e.g. that by Hagemann ("Allgemeine Genetik", Gustav Fischer Verlag, Stuttgart, 1986).

Possible mutations are transitions, transversions, insertions and deletions. Depending on the effect of the amino acid exchange on the enzyme activity, "missense mutations" or "nonsense mutations" are referred to. Insertions or deletions of at least one base pair in a gene lead to "frame shift mutations", as a consequence of which incorrect amino acids are incorporated or translation is interrupted prematurely. Deletions of several codons typically lead to a complete loss of the enzyme activity. Instructions on generation of such mutations are prior art and can be found in known textbooks of genetics and molecular biology, such as e.g. the textbook by Knippers ("Molekulare Genetik", 7th edition, Georg Thieme Verlag, Stuttgart, Germany, 1997), that by Winnacker ("Gene und Klone", VCH Verlagsgesellschaft, Weinheim, Germany, 1999) or that by Hagemann ("Allgemeine Genetik", 4th edition, Gustav Fischer Verlag, Stuttgart, 1999).

A common method of mutating genes of C. glutamicum is the method of "gene disruption" and "gene replacement" described by Schwarzer and Pühler (Bio/Technology 9, 84-87 (1991)).

In the method of gene disruption a central part of the coding region of the gene of interest is cloned in a plasmid vector which can replicate in a host (typically E. coli), but not in C. glutamicum. Possible vectors are, for example, pSUP301 (Simon et al., Bio/Technology 1, 784-791 (1983)), pK18mob or pK19mob (Schäfer et al., Gene 145, 69-73 (1994)), pK18mobsacB or pK19mobsacB (Jäger et al., Journal of Bacteriology 174: 5462-65 (1992)), pGEM-T (Promega corporation, Madison, WI, USA), pCR2.1-TOPO (Shuman (1994). Journal of Biological Chemistry 269:32678-84; US Patent 5,487,993), pCR®Blunt

(Invitrogen, Groningen, The Netherlands; Bernard et al.,  
Journal of Molecular Biology, 234: 534-541 (1993)) or pEM1  
(Schrumpf et al, 1991, Journal of Bacteriology 173:4510-4516).  
The plasmid vector which contains the central part of the  
5 coding region of the gene is then transferred into the desired  
strain of *C. glutamicum* by conjugation or transformation. The  
method of conjugation is described, for example, by Schäfer et  
al. (Applied and Environmental Microbiology 60, 756-759  
(1994)). Methods for transformation are described, for  
10 example, by Thierbach et al. (Applied Microbiology and  
Biotechnology 29, 356-362 (1988)), Dunican and Shivnan  
(Bio/Technology 7, 1067-1070 (1989)) and Tauch et al. (FEMS  
Microbiological Letters 123, 343-347 (1994)). After  
homologous recombination by means of a "cross over" event, the  
15 coding region of the gene in question is interrupted by the  
vector sequence and two incomplete alleles are obtained, one  
lacking the 3' end and one lacking the 5' end. This method  
has been used, for example, by Fitzpatrick et al. (Applied  
Microbiology and Biotechnology 42, 575-580 (1994)) to  
20 eliminate the *recA* gene of *C. glutamicum*.

In the method of "gene replacement", a mutation, such as e.g.  
a deletion, insertion or a base exchange, is established in  
vitro in the gene of interest. The allele prepared is in turn  
cloned in a vector which is not replicative for *C. glutamicum*  
and this is then transferred into the desired host of *C.*  
25 *glutamicum* by transformation or conjugation. After homologous  
recombination by means of a first "cross over" event which  
effects integration and a suitable second "cross-over" event  
which effects excision in the target gene or in the target  
30 sequence, the incorporation of the mutation or of the allele  
is achieved. This method was used, for example, by Peters-  
Wendisch et al. (Microbiology 144, 915 - 927 (1998)) to  
eliminate the *pyc* gene of *C. glutamicum* by a deletion.

A deletion, insertion or a base exchange can be incorporated  
35 into the *metR* gene or the *metZ* gene in this manner.

In addition, it may be advantageous for the production of L-amino acids, in particular L-methionine, to enhance, in particular to over-express, one or more enzymes of the particular biosynthesis pathway, of glycolysis, of anaplerosis, of the citric acid cycle or of amino acid export, in addition to attenuation of the metR gene and/or of the metZ gene.

The term "enhancement" in this connection describes the increase in the intracellular activity of one or more enzymes (proteins) in a microorganism which are coded by the corresponding DNA, for example by increasing the number of copies of the gene or genes, using a potent promoter or using a gene or allele which codes for a corresponding enzyme (protein) having a high activity, and optionally combining these measures.

By enhancement measures, in particular over-expression, the activity or concentration of the corresponding protein is in general increased by at least 10%, 25%, 50%, 75%, 100%, 150%, 200%, 300%, 400% or 500%, up to a maximum of 1000% or 2000%, based on the starting microorganism.

Thus, for the preparation of L-methionine, one or more genes chosen from the group consisting of

- the gap gene which codes for glyceraldehyde 3-phosphate dehydrogenase (Eikmanns (1992), Journal of Bacteriology 174:6076-6086),
- the tpi gene which codes for triose phosphate isomerase (Eikmanns (1992), Journal of Bacteriology 174:6076-6086),
- the pgk gene which codes for 3-phosphoglycerate kinase (Eikmanns (1992), Journal of Bacteriology 174:6076-6086),
- the pyc gene which codes for pyruvate carboxylase (Peters-Wendisch et al. (Microbiology 144, 915 - 927 (1998)),
- the lysC gene which codes for a feed-back resistant aspartate kinase (Accession No.P26512),

- the metA gene which codes for homoserine O-acetyltransferase (ACCESSION Number AF052652; EP-B-0387527; EP-A-0699759),
- the metB gene which codes for cystathionine gamma-synthase (ACCESSION Number AF126953),
- 5 • the aecD gene which codes for cystathionine gamma-lyase (ACCESSION Number M89931),
- the glyA gene which codes for serine hydroxymethyltransferase (JP-A-08107788),
- the metY gene which codes for O-acetylhomoserine
- 10 sulfhydrylase (DSM 13556)

can be enhanced, in particular over-expressed.

It may furthermore be advantageous for the production of L-methionine, in addition to the attenuation of the metR and/or metZ gene, for one or more genes chosen from the group

15 consisting of

- the thrB gene which codes for homoserine kinase (ACCESSION Number P08210),
- the ilvA gene which codes for threonine dehydratase (ACCESSION Number Q04513),
- 20 • the thrC gene which codes for threonine synthase (ACCESSION Number P23669),
- the ddh gene which codes for meso-diaminopimelate D-dehydrogenase (ACCESSION Number Y00151)
- the pck gene which codes for phosphoenol pyruvate
- 25 carboxykinase (DE 199 50 409.1; DSM 13047),
- the pgi gene which codes for glucose 6-phosphate isomerase (US 09/396,478; DSM 12969),
- the poxB gene which codes for pyruvate oxidase (DE: 1995 1975.7)

to be attenuated, in particular for the expression thereof to be reduced.

In addition to attenuation of the metR gene and/or of the metZ gene it may furthermore be advantageous, for the production of amino acids, in particular L-methionine, to eliminate undesirable side reactions (Nakayama: "Breeding of Amino Acid Producing Micro-organisms", in: Overproduction of Microbial Products, Krumphanzl, Sikyta, Vanek (eds.), Academic Press, London, UK, 1982).

The invention also provides the microorganisms prepared according to the invention, and these can be cultured continuously or discontinuously in the batch process (batch culture) or in the fed batch (feed process) or repeated fed batch process (repetitive feed process) for the purpose of production of L-amino acids, in particular L-methionine. A summary of known culture methods is described in the textbook by Chmiel (Bioprozesstechnik 1. Einführung in die Bioverfahrenstechnik (Gustav Fischer Verlag, Stuttgart, 1991)) or in the textbook by Storhas (Bioreaktoren und periphere Einrichtungen (Vieweg Verlag, Braunschweig/ Wiesbaden, 1994)).

The culture medium to be used must meet the requirements of the particular strains in a suitable manner. Descriptions of culture media for various microorganisms are contained in the handbook "Manual of Methods for General Bacteriology" of the American Society for Bacteriology (Washington D.C., USA, 1981).

Sugars and carbohydrates, such as e.g. glucose, sucrose, lactose, fructose, maltose, molasses, starch and cellulose, oils and fats, such as, for example, soya oil, sunflower oil, groundnut oil and coconut fat, fatty acids, such as, for example, palmitic acid, stearic acid and linoleic acid, alcohols, such as, for example, glycerol and ethanol, and organic acids, such as, for example, acetic acid, can be used as the source of carbon. These substances can be used individually or as a mixture.

Organic nitrogen-containing compounds, such as peptones, yeast extract, meat extract, malt extract, corn steep liquor, soya bean flour and urea, or inorganic compounds, such as ammonium sulfate, ammonium chloride, ammonium phosphate, ammonium carbonate and ammonium nitrate, can be used as the source of nitrogen. The sources of nitrogen can be used individually or as a mixture.

Organic and inorganic sulfur-containing compounds, such as, for example, sulfides, sulfites, sulfates and thiosulfates, can be used as a source of sulfur, in particular for the preparation of methionine.

Phosphoric acid, potassium dihydrogen phosphate or dipotassium hydrogen phosphate or the corresponding sodium-containing salts can be used as the source of phosphorus. The culture medium must furthermore comprise salts of metals, such as, for example, magnesium sulfate or iron sulfate, which are necessary for growth. Finally, essential growth substances, such as amino acids and vitamins, can be employed in addition to the above-mentioned substances. Suitable precursors can moreover be added to the culture medium. The starting substances mentioned can be added to the culture in the form of a single batch, or can be fed in during the culture in a suitable manner.

Basic compounds, such as sodium hydroxide, potassium hydroxide, ammonia or aqueous ammonia, or acid compounds, such as phosphoric acid or sulfuric acid, can be employed in a suitable manner to control the pH of the culture. Antifoams, such as, for example, fatty acid polyglycol esters, can be employed to control the development of foam. Suitable substances having a selective action, such as, for example, antibiotics, can be added to the medium to maintain the stability of plasmids. To maintain aerobic conditions, oxygen or oxygen-containing gas mixtures, such as, for example, air, are introduced into the culture. The temperature of the culture is usually 20°C to 45°C, and preferably 25°C to 40°C. Culturing is continued until a maximum of the desired product



has formed. This target is usually reached within 10 hours to 160 hours.

The fermentation broths obtained in this way, in particular containing L-methionine, usually have a dry weight of 7.5 to 25 wt.% and contain L-methionine. It is furthermore also advantageous if the fermentation is conducted in a sugar-limited procedure at least at the end, but in particular over at least 30% of the duration of the fermentation. That is to say, the concentration of utilizable sugar in the fermentation medium is reduced to  $\geq 0$  to 3 g/l during this period.

The fermentation broth prepared in this manner, in particular containing L-methionine, is then further processed. Depending on requirements, the all or some of the biomass can be removed from the fermentation broth by separation methods, such as centrifugation, filtration, decanting or a combination thereof, or it can be left completely in. This broth is then thickened or concentrated by known methods, such as with the aid of a rotary evaporator, thin film evaporator, falling film evaporator, by reverse osmosis, or by nanofiltration. This concentrated fermentation broth can then be worked up by methods of freeze drying, spray drying, spray granulation or by other processes to give a preferably free-flowing, finely divided powder.

This free-flowing, finely divided powder can then in turn be converted by suitable compacting or granulating processes into a coarse-grained, readily free-flowing, storable and largely dust-free product. In the granulation or compacting it is advantageous to employ conventional organic or inorganic auxiliary substances or carriers, such as starch, gelatin, cellulose derivatives or similar substances, such as are conventionally used as binders, gelling agents or thickeners in foodstuffs or feedstuffs processing, or further substances, such as, for example, silicas, silicates or stearates.

"Free-flowing" is understood as meaning powders which flow unimpeded out of the vessel with the opening of 5 mm (millimeters) of a series of glass outflow vessels with

outflow openings of various sizes (Klein, Seifen, Öle, Fette, Wachse 94, 12 (1968)).

As described here, "finely divided" means a powder with a predominant content (> 50 %) having a particle size of 20 to 200 µm diameter. "Coarse-grained" means products with a predominant content (> 50 %) having a particle size of 200 to 2000 µm diameter. In this context, "dust-free" means that the product contains only small contents (< 5 %) having particle sizes of less than 20 µm diameter. The particle size determination can be carried out with methods of laser diffraction spectrometry. The corresponding methods are described in the textbook on "Teilchengrößenmessung in der Laborpraxis" by R. H. Müller and R. Schuhmann, Wissenschaftliche Verlagsgesellschaft Stuttgart (1996) or in the textbook "Introduction to Particle Technology" by M. Rhodes, Verlag Wiley & Sons (1998).

"Storable" in the context of this invention means a product which can be stored for up to 120 days, preferably up to 52 weeks, particularly preferably 60 months, without a substantial loss (< 5%) of methionine occurring.

Alternatively, however, the product can be absorbed on to an organic or inorganic carrier substance which is known and conventional in feedstuffs processing, for example, silicas, silicates, grits, brans, meals, starches, sugars or others, and/or mixed and stabilized with conventional thickeners or binders. Use examples and processes in this context are described in the literature (Die Mühle + Mischfuttertechnik 132 (1995) 49, page 817).

Finally, the product can be brought into a state in which it is stable to digestion by animal stomachs, in particular the stomach of ruminants, by coating processes ("coating") using film-forming agents, such as, for example, metal carbonates, silicas, silicates, alginates, stearates, starches, gums and cellulose ethers, as described in DE-C-4100920.

If the biomass is separated off during the process, further inorganic solids, for example added during the fermentation, are in general removed. In addition, the animal feedstuffs additive according to the invention comprises at least the predominant proportion of the further substances, in particular organic substances, which are formed or added and are present in solution in the fermentation broth, where these have not been separated off by suitable processes.

In one aspect of the invention, the biomass can be separated off to the extent of up to 70%, preferably up to 80%, preferably up to 90%, preferably up to 95%, and particularly preferably up to 100%. In another aspect of the invention, up to 20% of the biomass, preferably up to 15%, preferably up to 10%, preferably up to 5%, particularly preferably no biomass is separated off.

These organic substances include organic by-products which are optionally produced, in addition to the L-methionine, and optionally discharged by the microorganisms employed in the fermentation. These include L-amino acids chosen from the group consisting of L-lysine, L-valine, L-threonine, L-alanine or L-tryptophan. They include vitamins chosen from the group consisting of vitamin B1 (thiamine), vitamin B2 (riboflavin), vitamin B5 (pantothenic acid), vitamin B6 (pyridoxine), vitamin B12 (cyanocobalamin), nicotinic acid/nicotinamide and vitamin E (tocopherol). They also include organic acids which carry one to three carboxyl groups, such as, acetic acid, lactic acid, citric acid, malic acid or fumaric acid. Finally, they also include sugars, for example, trehalose. These compounds are optionally desired if they improve the nutritional value of the product.

These organic substances, including L-methionine and/or D-methionine and/or the racemic mixture D,L-methionine, can also be added, depending on requirements, as a concentrate or pure substance in solid or liquid form during a suitable process step. These organic substances mentioned can be added individually or as mixtures to the resulting or concentrated

fermentation broth, or also during the drying or granulation process. It is likewise possible to add an organic substance or a mixture of several organic substances to the fermentation broth and a further organic substance or a further mixture of several organic substances during a later process step, for example granulation.

The product described above is suitable as a feedstuffs additive, i.e. feed additive, for animal nutrition.

10 The L-methionine content of the animal feedstuffs additive is conventionally 1 wt.% to 80 wt.%, preferably 2 wt.% to 80 wt.%, particularly preferably 4 wt.% to 80 wt.%, and very particularly preferably 8 wt.% to 80 wt.%, based on the dry weight of the animal feedstuffs additive. Contents of 1 wt.% to 60 wt.%, 2 wt.% to 60 wt.%, 4 wt.% to 60 wt.%, 6 wt.% to 60 wt.%, 1 wt.% to 40 wt.%, 2 wt.% to 40 wt.% or 4 wt.% to 40 wt.% are likewise possible. The water content of the feedstuffs additive is conventionally up to 5 wt.%, preferably up to 4 wt.%, and particularly preferably less than 2 wt.%.

20 The invention also provides a process for the preparation of an L-methionine-containing animal feedstuffs additive from fermentation broths, which comprises the steps

- a) culture and fermentation of an L-methionine-producing microorganism in a fermentation medium;
- b) removal of water from the L-methionine-containing fermentation broth (concentration);
- c) removal of an amount of 0 to 100 wt.% of the biomass formed during the fermentation; and
- d) drying of the fermentation broth obtained according to a) and/or b) to obtain the animal feedstuffs additive in the desired powder or granule form.

If desired, one or more of the following steps can furthermore be carried out in the process according to the invention:



## Example 1

Preparation of a genomic cosmid gene library from  
*Corynebacterium glutamicum* ATCC 13032

Chromosomal DNA from *Corynebacterium glutamicum* ATCC 13032 was  
isolated as described by Tauch et al. (1995, Plasmid 33:168-  
179) and partly cleaved with the restriction enzyme Sau3AI  
(Amersham Pharmacia, Freiburg, Germany, Product Description  
Sau3AI, Code no. 27-0913-02). The DNA fragments were  
dephosphorylated with shrimp alkaline phosphatase (Roche  
Diagnostics GmbH, Mannheim, Germany, Product Description SAP,  
Code no. 1758250). The DNA of the cosmid vector SuperCos1  
(Wahl et al. (1987) Proceedings of the National Academy of  
Sciences USA 84:2160-2164), obtained from Stratagene (La  
Jolla, USA, Product Description SuperCos1 Cosmid Vector Kit,  
Code no. 251301) was cleaved with the restriction enzyme XbaI  
(Amersham Pharmacia, Freiburg, Germany, Product Description  
XbaI, Code no. 27-0948-02) and likewise dephosphorylated with  
shrimp alkaline phosphatase.

The cosmid DNA was then cleaved with the restriction enzyme  
BamHI (Amersham Pharmacia, Freiburg, Germany, Product  
Description BamHI, Code no. 27-0868-04). The cosmid DNA  
treated in this manner was mixed with the treated ATCC13032  
DNA and the batch was treated with T4 DNA ligase (Amersham  
Pharmacia, Freiburg, Germany, Product Description T4-DNA-  
Ligase, Code no. 27-0870-04). The ligation mixture was then  
packed in phages with the aid of Gigapack II XL Packing  
Extract (Stratagene, La Jolla, USA, Product Description  
Gigapack II XL Packing Extract, Code no. 200217).

For infection of the *E. coli* strain NM554 (Raleigh et al.  
1988, Nucleic Acid Research 16:1563-1575) the cells were taken  
up in 10 mM MgSO<sub>4</sub> and mixed with an aliquot of the phage  
suspension. The infection and titering of the cosmid library  
were carried out as described by Sambrook et al. (1989,  
Molecular Cloning: A Laboratory Manual, Cold Spring Harbor),  
the cells being plated out on LB agar (Lennox, 1955, Virology,

1:190) with 100 mg/l ampicillin. After incubation overnight at 37°C, recombinant individual clones were selected.

## Example 2

### Isolation and sequencing of the metR and metZ genes

5 The cosmid DNA of an individual colony was isolated with the Qiaprep Spin Miniprep Kit (Product No. 27106, Qiagen, Hilden, Germany) in accordance with the manufacturer's instructions and partly cleaved with the restriction enzyme Sau3AI (Amersham Pharmacia, Freiburg, Germany, Product Description  
10 Sau3AI, Product No. 27-0913-02). The DNA fragments were dephosphorylated with shrimp alkaline phosphatase (Roche Diagnostics GmbH, Mannheim, Germany, Product Description SAP, Product No. 1758250). After separation by gel  
15 electrophoresis, the cosmid fragments in the size range of 1500 to 2000 bp were isolated with the QiaExII Gel Extraction Kit (Product No. 20021, Qiagen, Hilden, Germany).

The DNA of the sequencing vector pZero-1, obtained from Invitrogen (Groningen, The Netherlands, Product Description Zero Background Cloning Kit, Product No. K2500-01) was cleaved  
20 with the restriction enzyme BamHI (Amersham Pharmacia, Freiburg, Germany, Product Description BamHI, Product No. 27-0868-04). The ligation of the cosmid fragments in the sequencing vector pZero-1 was carried out as described by Sambrook et al. (1989, Molecular Cloning: A laboratory Manual,  
25 Cold Spring Harbor), the DNA mixture being incubated overnight with T4 ligase (Pharmacia Biotech, Freiburg, Germany). This ligation mixture was then electroporated (Tauch et al. 1994, FEMS Microbiol Letters, 123:343-7) into the E. coli strain DH5αMCR (Grant, 1990, Proceedings of the National Academy of  
30 Sciences U.S.A., 87:4645-4649) and plated out on LB agar (Lennox, 1955, Virology, 1:190) with 50 mg/l zeocin.

The plasmid preparation of the recombinant clones was carried out with Biorobot 9600 (Product No. 900200, Qiagen, Hilden, Germany). The sequencing was carried out by the dideoxy chain  
35 termination method of Sanger et al. (1977, Proceedings of the

1. . .  
National Academy of Sciences U.S.A., 74:5463-5467) with  
modifications according to Zimmermann et al. (1990, Nucleic  
Acids Research, 18:1067). The "RR dRhodamin Terminator Cycle  
Sequencing Kit" from PE Applied Biosystems (Product No.  
5 403044, Weiterstadt, Germany) was used. The separation by gel  
electrophoresis and analysis of the sequencing reaction were  
carried out in a "Rotiphoresis NF Acrylamide/Bisacrylamide"  
Gel (29:1) (Product No. A124.1, Roth, Karlsruhe, Germany) with  
the "ABI Prism 377" sequencer from PE Applied Biosystems  
10 (Weiterstadt, Germany).

The raw sequence data obtained were then processed using the  
Staden program package (1986, Nucleic Acids Research, 14:217-  
231) version 97-0. The individual sequences of the pZero1  
derivatives were assembled to a continuous contig. The  
15 computer-assisted coding region analysis was prepared with the  
XNIP program (Staden, 1986, Nucleic Acids Research, 14:217-  
231).

The resulting nucleotide sequence is shown in SEQ ID No. 1.  
Analysis of the nucleotide sequence showed two open reading  
20 frames of 567 base pairs and 1146 base pairs, which were  
called the metR gene and metZ gene. The metR gene codes for a  
protein of 189 amino acids, the metZ gene codes for a protein  
of 382 amino acids.

### Example 3

25 Deletion of the metR and metZ genes in *C. glutamicum* ATCC13032

For this, chromosomal DNA was isolated from the strain  
ATCC13032 by the method of Tauch et al. (Plasmid 33:168-179  
(1995)). On the basis of the sequences of the metR and metZ  
genes known for *C. glutamicum* from example 2, the  
30 oligonucleotides described below were chosen for generation of  
the metR-metZ deletion allele by means of the polymerase chain  
reaction (PCR) by the gene Soeing method (Horton, Molecular  
Biotechnology 3: 93-98 (1995)).



orfR 20 (see also SEQ ID No. 4):

5'-GAT CTA GAA TTC AAC ATC GTC CGA TAG GCC AG-3'

orfRmetZ del (see also SEQ ID No. 5):

5'-GGT CTG GTT ACT GGC TAC GAA TGC AGA CAA TAA TGT CGC T-3'

5 metZ 21 (see also SEQ ID No. 6):

5'-GAT CTA GAA TTC CGC TTC CGA GAA TCA ATA CC-3'

The primers shown were synthesized by MWG Biotech (Ebersberg, Germany) and the PCR reaction was carried out using Pfu polymerase (Stratagene, Product. No. 600135, La Jolla, USA) and a PTC 100 Thermocycler (MJ Research Inc., Waltham, USA).

The primers orfR 20 and metZ 21 contain in each case an inserted cleavage site for the restriction enzyme EcoRI, which are marked by underlining in the nucleotide sequence shown above. The primer orfRmetZ del is composed of two regions of the nucleotide sequence, one of which bonds in the "upstream" region of metR and includes the first two nucleotides of the start codon ATG, and the other bonds after the stop codon of metZ in the "downstream" region thereof.

In a first PCR reaction, by means of the primers orfR 20 and orfRmetZ del an amplification product 402 bp in size was produced which contains the "upstream" region of the met R gene including the first two nucleotides of the start codon ATG, and additionally a 20 bp extension, appended with the oligonucleotide orfRmetZ del, which corresponds to a part of the nucleotide sequence of the "downstream" region of metZ. The amplification product was isolated from the agarose gel with the QiaExII Gel Extraction Kit (Product No. 20021, Qiagen, Hilden, Germany). It was called metRmetZ del fragment 1 and is shown in SEQ ID No. 7.

In a second PCR reaction the metRmetZ deletion derivative 982 bp in size was produced with the primer metZ 21 and the purified amplification product metRmetZ del fragment 1, which can bond by means of the 20 bp extension from the oligonucleotide orfRmetZ del in the "downstream" region of

metZ to the DNA and function there as a primer. It comprises, between the two EcoRI cleavage sites appended to the primers, 368 bp of the "upstream region" of the metR gene, the first two nucleotides of the start codon ATG and 588 bp of the "downstream" region of the metZ gene, starting with the fourth nucleotide after the stop codon of the metZ gene.

It was called metRmetZ del fragment 2 and is shown in SEQ ID No. 8.

#### Example 4

10 Deletion mutagenesis of the metR and metZ genes in *C. glutamicum* ATCC13032

The 982 bp metRmetZ deletion derivative obtained in example 3 was incorporated by means of deletion mutagenesis with the aid of the sacB system described by Schäfer et al., Gene, 14, 69-73 (1994) into the chromosome of *C. glutamicum*. This system enables the expert to identify or select allele exchanges which take place by homologous recombination.

##### 1. Construction of the exchange vector

###### pK18mobsacBdeltametRmetZ

20 The metRmetZ deletion derivative 982 bp in size obtained in example 3 was cleaved with the restriction endonuclease EcoRI and then isolated from the agarose gel with the QiaExII Gel Extraction Kit (Product No. 20021, Qiagen, Hilden, Germany) and used for ligation with the mobilizable cloning vector pK18mobsacB described by Schäfer et al., Gene, 14, 69-73 (1994). This was cleaved beforehand with the restriction enzyme EcoRI, subsequently dephosphorylated with shrimp alkaline phosphatase (Roche Diagnostics GmbH, Mannheim, Germany, Product Description SAP, Product No. 1758250) and then mixed with the metRmetZ deletion derivative, and the mixture was treated with T4 DNA ligase (Amersham-Pharmacia, Freiburg, Germany).

The *E. coli* strain S17-1 (Simon et al., Bio/Technologie [Bio/Technology] 1: 784-791, 1993) was then electroporated

with the ligation batch (Hanahan, In. DNA cloning. A practical approach. Vol.1. ILR-Press, Cold Spring Harbor, New York, 1989). Selection of plasmid-carrying cells was made by plating out the transformation batch on LB agar (Sambrock et al.,  
5 Molecular Cloning: A Laboratory Manual, 2<sup>nd</sup> Ed., Cold Spring Harbor, New York, 1989), which had been supplemented with 25 mg/l kanamycin.

Plasmid DNA was isolated from a transformant with the aid of the QIAprep Spin Miniprep Kit from Qiagen and the cloned lrp  
10 deletion allele was verified by means of sequencing by MWG Biotech (Ebersberg, Germany).

The plasmid was called pK18mobsacBdeltametRmetZ and is shown in figure 1.

#### Example 5

15 Deletion mutagenesis of the metR and metZ genes in the C. glutamicum strain ATCC13032

The vector pK18mobsacBdeltametRmetZ mentioned in example 4 was electroporated by the electroporation method of Tauch et al., (1989 FEMS Microbiology Letters 123: 343-347) in the  
20 strain C. glutamicum ATCC13032. The vector cannot replicate independently in ATCC13032 and is retained in the cell only if it has integrated into the chromosome. Selection of clones with integrated pK18mobsacBdeltametRmetZ was carried out by plating out the electroporation batch on LB agar (Sambrock et  
25 al., Molecular Cloning: A Laboratory Manual, 2<sup>nd</sup> Ed., Cold Spring Harbor, New York, 1989), which had been supplemented with 15 mg/l kanamycin. Clones which had grown on were plated out on LB agar plates with 25 mg/l kanamycin and incubated for 16 hours at 33°C. To achieve excision of the plasmid together  
30 with the complete chromosomal copy of the metR and metZ genes, the clones were incubated unselectively overnight in LB medium and then cultured on LB agar with 10% sucrose.

The plasmid pK18mobsacB contains a copy of the sacB gene, which converts sucrose into levan sucrose, which is toxic to

C. glutamicum. Only those clones in which the pK18mobsacBdeltametRmetZ integrated has been excised again therefore grow on LB agar with sucrose. In the excision, together with the plasmid either the complete chromosomal copy of the metR and metZ genes can be excised, or the metRmetZ deletion derivative. To demonstrate that the metR and metZ genes are deleted in the chromosome, the plasmid pK18mobsacBdeltametRmetZ was marked by the method of "The DIG System Users Guide for Filter Hybridization" of Boehringer Mannheim GmbH (Mannheim, Germany, 1993) using the DIG hybridization kit from Boehringer. Chromosomal DNA of a potential deletion mutant was isolated by the method of Eikmanns et al. (Microbiology 140: 1817-1828 (1994)) and in each case cleaved with the restriction enzymes HindIII and EcoRI in separate batches.

The fragments formed were separated by agarose gel electrophoresis and hybridized at 68°C with the Dig hybridization kit from Boehringer. With the aid of the fragments formed, it could be shown that the strain ATCC13032 has lost its copies of the metR and metZ genes, and instead the region from the 3rd nucleotide of the metR gene up to and including the 3rd nucleotide after the metZ gene is deleted.

The strain was called C. glutamicum ATCC13032deltametRmetZ.

#### Example 6

Preparation of methionine with the strain C. glutamicum ATCC13032deltametRmetZ

The C. glutamicum strain ATCC13032deltametRmetZ obtained in example 4 was cultured in a nutrient medium suitable for the production of methionine and the methionine content in the culture supernatant was determined.

For this, the strain was first incubated on a brain-heart agar plate for 24 hours at 33°C. Starting from this agar plate culture, a preculture was seeded (10 ml medium in a 100 ml

conical flask). The medium MM was used as the medium for the preculture.

#### Medium MM

CSL (corn steep liquor)	5 g/l
MOPS (morpholinopropanesulfonic acid)	20 g/l
Glucose (autoclaved separately)	50g/l
Salts:	
(NH <sub>4</sub> ) <sub>2</sub> SO <sub>4</sub>	25 g/l
KH <sub>2</sub> PO <sub>4</sub>	0.1 g/l
MgSO <sub>4</sub> * 7 H <sub>2</sub> O	1.0 g/l
CaCl <sub>2</sub> * 2 H <sub>2</sub> O	10 mg/l
FeSO <sub>4</sub> * 7 H <sub>2</sub> O	10 mg/l
MnSO <sub>4</sub> * H <sub>2</sub> O	5.0mg/l
Biotin (sterile-filtered)	0.01 mg/l
Vitamin B12 (sterile-filtered)	0.02 mg/l
Thiamine * HCl (sterile-filtered)	0.2 mg/l
CaCO <sub>3</sub>	25 g/l

- 5 The CSL, MOPS and the salt solution were brought to pH 7 with aqueous ammonia and autoclaved. The sterile substrate and vitamin solutions were then added, as well as the CaCO<sub>3</sub> autoclaved in the dry state.

- 10 The preculture was incubated for 16 hours at 33°C at 240 rpm on a shaking machine. A main culture was seeded from this preculture such that the initial OD (660 nm) of the main culture was 0.1. Medium MM was also used for the main culture.

Culturing is carried out in a 10 ml volume in a 100 ml conical flask with baffles. Culturing was carried out at 33°C and 80% atmospheric humidity.

After 72 hours, the OD was determined at a measurement wavelength of 660 nm with a Biomek 1000 (Beckmann Instruments GmbH, Munich). The amount of methionine formed was determined with an amino acid analyzer from Eppendorf-BioTronik (Hamburg, Germany) by ion exchange chromatography and post-column derivation with ninhydrin detection.

10 The result of the experiment is shown in table 1.

Table 1

Strain	OD (660 nm)	Methionine mg/l
ATCC13032	11.3	1.5
ATCC13032deltametRmetZ	12.0	9.4

Brief Description of the Figure:

Figure 1: Map of the plasmid pK18mobsacBdeltametRmetZ.

15 The abbreviations and designations used have the following meaning:

sacB                      sacB gene

oriV                      ColE1-similar origin from pMB1

KmR                      Kanamycin resistance

HindIII                  Cleavage site of the restriction enzyme  
HindIII

EcoRI                    Cleavage site of the restriction enzyme EcoRI

RP4mob            RP4 mobilization site

orfR'            Nucleotides A and T of the start codon of  
                 metR

metRmetZ del    Cloned deletion derivative for metR and metZ

This disclosure is based on priority documents DE 100 43  
335.9, DE 101 09 688.7 and US 60/294,224, each incorporated by  
reference.

- 5    Obviously, numerous modifications of the invention are  
possible in view of the above teachings. Therefore, within  
the scope of the appended claims, the invention may be  
practiced otherwise than as specifically described herein.

SEQUENCE PROTOCOL

<110> Degussa AG

5 <120> Nucleotide sequences which code for the metR and metZ genes

<130> 000369 BT

<140>

10 <141>

<160> 8

<170> PatentIn Ver. 2.1

15

<210> 1

<211> 2628

<212> DNA

<213> Corynebacterium glutamicum

20

<220>

<221> CDS

<222> (447)..(1013)

<223> metR gene

25

<220>

<221> CDS

<222> (1038)..(2183)

<223> metZ gene

30

<400> 1

cggtcacgtt gggatcgttg tcaaaactcc ccagtgggtt cacttcataa actcgcgagg 60

ttttccgggg aactgaaaaa catcgctcca taggccagcg tctaattcag cagcgatttc 120

ggcagcaagc ccagcgccat taatcagagc ggtgaaataa acatgggttca tgattatgtc 180

aggacggtaa ttagacttat gaccagggtt aaggagggtca ccagggttgaa gccgcgctat 240

35

tgttccgtgg aaaagggggc cctgatctag ctgattattc atcgcagtaa gcgctttcgg 300

taggtgggtg aatcatcgta gtcttccgag ccccgtagacc cgatccgttt tgtgcaatcc 360

aatgctactc ccacagagcg ggctactttc tctaaaaatg ttctcatagt agataaaatt 420

45

gttcttaaag cgacattatt gtctgc atg gaa gac gat ctc agt gct gct ctc 473

Met Glu Asp Asp Leu Ser Ala Ala Leu

1

5

gtc aaa gcg ctg ttc gac gcg cga acc caa cgc agg ctc tct atc tcg 521

Val Lys Ala Leu Phe Asp Ala Arg Thr Gln Arg Arg Leu Ser Ile Ser

10

15

20

25

gcg tta gct gaa tcc tcc ggt gtg tcg cga gca atg att tcc cgc gtg 569

Ala Leu Ala Glu Ser Ser Gly Val Ser Arg Ala Met Ile Ser Arg Val

30

35

40



5	ggt gca ttg ggt atg acg ctt tcg gag ctc att gca cag gct gaa ggt	665
	Gly Ala Leu Gly Met Thr Leu Ser Glu Leu Ile Ala Gln Ala Glu Gly	
	60 65 70	
10	ggc tat gac cgg ggc gct cgg cgg tca aag cag tct gta tgg aca gat	713
	Gly Tyr Asp Arg Gly Ala Arg Arg Ser Lys Gln Ser Val Trp Thr Asp	
	75 80 85	
15	cca gct acc ggt tac aca cgg cgt gca gtg tca cag ccg tca gaa tcc	761
	Pro Ala Thr Gly Tyr Thr Arg Arg Ala Val Ser Gln Pro Ser Glu Ser	
	90 95 100 105	
20	cca cta gaa cta gtg gaa gta atg ctg cct cct ggg gcg gaa gtt ggc	809
	Pro Leu Glu Leu Val Glu Val Met Leu Pro Pro Gly Ala Glu Val Gly	
	110 115 120	
25	tac cca gct gat gct tat cgt ttc atg gat cag gtg gtc tgg gta ctc	857
	Tyr Pro Ala Asp Ala Tyr Arg Phe Met Asp Gln Val Val Trp Val Leu	
	125 130 135	
30	gaa ggg gcc gtt cgt att act gaa ggt gaa gag gtc cac gaa ctt tca	905
	Glu Gly Ala Val Arg Ile Thr Glu Gly Glu Glu Val His Glu Leu Ser	
	140 145 150	
35	acg ggg gat tgt cta cgg ttt ggg cct ccg cga gat acc gac ttt gct	953
	Thr Gly Asp Cys Leu Arg Phe Gly Pro Pro Arg Asp Thr Asp Phe Ala	
	155 160 165	
40	aat ccc acc acc gta gcc act agg tat tta gtt gcc ttg gac aag cgt	1001
	Asn Pro Thr Thr Val Ala Thr Arg Tyr Leu Val Ala Leu Asp Lys Arg	
	170 175 180 185	
45	gta cct cgt gct tgatataaca agtaaggaag cctg atg aat ttt tac cca	1052
	Val Pro Arg Ala Met Asn Phe Tyr Pro	
	190	
50	cca tct gta cct att aac cct gcg tgg cgt cca ccc aca gta act gtg	1100
	Pro Ser Val Pro Ile Asn Pro Ala Trp Arg Pro Pro Thr Val Thr Val	
	195 200 205 210	
55	caa gcg gga cgg cca gcc aga act cct ggt gcg ccg atg aac cca cct	1148
	Gln Ala Gly Arg Pro Ala Arg Thr Pro Gly Ala Pro Met Asn Pro Pro	
	215 220 225	
60	atc acg ttg tcc agc act tat gtt cat gat tca gaa aaa gct tat ggg	1196
	Ile Thr Leu Ser Thr Tyr Val His Asp Ser Glu Lys Ala Tyr Gly	
	230 235 240	
65	cgc gat ggc aat gat gga tgg ggt gca ttt gag gct gcc atg gga act	1244
	Arg Asp Gly Asn Asp Gly Trp Gly Ala Phe Glu Ala Ala Met Gly Thr	
	245 250 255	
70	cta gat ggt ggg ttc gcg gta tct tat tct tca ggt ttg gca gcg gca	1292
	Leu Asp Gly Gly Phe Ala Val Ser Tyr Ser Ser Gly Leu Ala Ala Ala	
	260 265 270	
75	acg tcg att gct gat ttg gtt cct act ggt ggc aca gtt gtt tta cct	1340

	Thr 275	Ser	Ile	Ala	Asp	Leu 280	Val	Pro	Thr	Gly	Gly 285	Thr	Val	Val	Leu	Pro 290	
5	aaa Lys	gct Ala	gcc Ala	tat Tyr	tat Tyr 295	ggc Gly	gtg Val	acc Thr	aat Asn	att Ile 300	ttc Phe	gcc Ala	agg Arg	atg Met	gaa Glu 305	gcc Ala	1388
10	cgc Arg	gga Gly	agg Arg	ctg Leu 310	aag Lys	gtt Val	cga Arg	act Thr	gtt Val 315	gat Asp	gca Ala	gac Asp	aat Asn	acc Thr 320	gaa Glu	gaa Glu	1436
15	gtg Val	att Ile	gct Ala	gct Ala	gct Ala	caa Gln	ggc Gly	gca Ala	gat Asp	gtg Val	gtg Val	tgg Trp	gtg Val	gaa Glu	tgc Ser	atc Ile	1484
20	gct Ala	aat Asn 340	ccg Pro	acg Thr	atg Met	gtg Val	gta Val	gct Ala	gat Asp	atc Ile	cct Pro	gca Ala	ata Ile	gtc Val	gac Asp	ggc Gly	1532
25	gtg Val 355	cgt Arg	ggg Gly	ctt Leu	gga Gly	gtt Val	ttg Leu	act Thr	gtc Val	gtt Val	gac Asp 365	gcg Ala	act Thr	ttc Phe	gca Ala	acg Thr 370	1580
30	cca Pro	ctt Leu	cgt Arg	caa Gln	cgt Arg 375	cca Pro	ttg Leu	gaa Glu	ctt Leu	ggc Gly 380	gct Ala	gat Asp	att Ile	gtg Val	ctt Leu 385	tac Tyr	1628
35	tgc Ser	gca Ala	acc Thr	aaa Lys 390	ctt Leu	atc Ile	ggc Gly	gga Gly	cac His 395	tct Ser	gat Asp	ctt Leu	ctt Leu	ctt Leu 400	gga Gly	gtc Val	1676
40	gca Ala	gtg Val	tgc Cys 405	aag Lys	tct Ser	gag Glu	cac His	cat His 410	gcg Ala	cag Gln	ttt Phe	ctt Leu	gcc Ala	act Thr 415	cac His	cgt Arg	1724
45	cat His	gat Asp 420	cat His	ggc Gly	tca Ser	gtg Val	ccg Pro	gga Gly	ggc Gly	ctt Leu	gaa Glu	gcg Ala	ttt Phe	ctt Leu 430	gct Ala	ctc Leu	1772
50	cgt Arg 435	gga Gly	ttg Leu	tat Tyr	tcc Ser	ttg Leu	gcg Ala	gtg Val	cgt Arg	ctt Leu	gat Asp 445	cga Arg	gca Ala	gaa Glu	tcc Ser	aac Asn 450	1820
	gca Ala	gca Ala	gaa Glu	ctt Leu	tgc Ser	ccg Arg	cga Arg	ctt Leu	aac Asn	gcg Ala 460	cat His	cct Pro	tgc Ser	gtt Val	acc Thr 465	cgc Arg	1868
	gtc Val	aat Asn	tat Tyr	cca Pro	gga Gly	ctt Leu	cct Pro	gat Asp	gat Asp 475	ccc Pro	caa Gln	cat His	gaa Glu	aaa Lys 480	gcc Ala	gtg Val	1916
	cga Arg	gtc Val	cta Leu	ccc Pro	tct Ser	gga Gly	tgt Cys	gga Gly	aac Asn 490	atg Met	ttg Leu	tca Ser	ttt Phe	gag Glu	ctt Leu	gat Asp	1964

	gca	aca	cct	gaa	cga	act	gat	gag	att	ctc	gaa	agc	ctg	tca	ctt	tta	2012
	Ala	Thr	Pro	Glu	Arg	Thr	Asp	Glu	Ile	Leu	Glu	Ser	Leu	Ser	Leu	Leu	
		500					505					510					
5	acc	cac	gcg	acc	agt	tgg	gga	ggt	gtg	gaa	aca	gcc	att	gaa	cgt	cgc	2060
	Thr	His	Ala	Thr	Ser	Trp	Gly	Gly	Val	Glu	Thr	Ala	Ile	Glu	Arg	Arg	
		515				520					525					530	
	acc	agg	cgg	gat	gct	gaa	gtg	gtg	gca	gga	gta	ccg	atg	act	ctt	tgc	2108
10	Thr	Arg	Arg	Asp	Ala	Glu	Val	Val	Ala	Gly	Val	Pro	Met	Thr	Leu	Cys	
					535					540						545	
	cgc	gtt	tcc	gta	gga	att	gaa	gac	gtt	gaa	gat	cta	tgg	gaa	gac	ctc	2156
15	Arg	Val	Ser	Val	Gly	Ile	Glu	Asp	Val	Glu	Asp	Leu	Trp	Glu	Asp	Leu	
				550					555						560		
	aac	gcc	tca	atc	gac	aaa	gtt	ctg	ggc	tagaactcgt	agccagtaac						2203
	Asn	Ala	Ser	Ile	Asp	Lys	Val	Leu	Gly								
		565						570									
20	cagac	cttca	gtg	tttg	ggtt	gcc	actcc	cag	tgct	gggggc	acat	gatcag	cga	agttctt			2263
	caggat	cgc	gctt	gatct	caac	acccat	ttg	ggttggg	gcg	gtgagca	tca	aggtgtc					2323
25	ggctt	ccatc	acag	ctttgt	cttc	ccttgag	ctg	ggtcgatg	agtt	catcgg	gact	tcgggc					2383
	gtagct	gcga	ccga	acgtgg	atcg	ggtatc	atcc	aggatt	cct	acttgg	cacc	gccttg					2443
	tccct	gaagt	ccgaaa	agct	cac	ggtcg	gtc	ggtgacg	atc	gggaaga	tgg	acctgga					2503
30	gacag	acaca	cgtggg	gtcc	aatc	gtgtcc	ggctt	cttttc	caag	cttggc	ggt	agaacgc					2563
	gattt	gatcg	gcttgc	agat	cccc	gaagga	ttgg	ccggtg	gctt	cggcga	cgagg	gtgga					2623
35	gctca																2628
	<210>	2															
	<211>	189															
40	<212>	PRT															
	<213>	Corynebacterium glutamicum															
	<400>	2															
45	Met	Glu	Asp	Asp	Leu	Ser	Ala	Ala	Leu	Val	Lys	Ala	Leu	Phe	Asp	Ala	
	1				5					10					15		
	Arg	Thr	Gln	Arg	Arg	Leu	Ser	Ile	Ser	Ala	Leu	Ala	Glu	Ser	Ser	Gly	
				20					25					30			
	Val	Ser	Arg	Ala	Met	Ile	Ser	Arg	Val	Glu	Asn	Ala	Glu	Ala	Gln	Pro	
				35				40					45				
50	Ser	Ala	Ala	Leu	Leu	Gly	Arg	Leu	Ser	Gly	Ala	Leu	Gly	Met	Thr	Leu	
		50				55						60					
	Ser	Glu	Leu	Ile	Ala	Gln	Ala	Glu	Gly	Gly	Tyr	Asp	Arg	Gly	Ala	Arg	
		65				70					75					80	

	Glu	Gly	Glu	Glu	Val	His	Glu	Leu	Ser	Thr	Gly	Asp	Cys	Leu	Arg	Phe
	145					150					155					160
	Gly	Pro	Pro	Arg	Asp	Thr	Asp	Phe	Ala	Asn	Pro	Thr	Thr	Val	Ala	Thr
					165					170						175
5	Arg	Tyr	Leu	Val	Ala	Leu	Asp	Lys	Arg	Val	Pro	Arg	Ala			
				180					185							
	<210> 3															
10	<211> 382															
	<212> PRT															
	<213> Corynebacterium glutamicum															
	<400> 3															
15	Met	Asn	Phe	Tyr	Pro	Pro	Ser	Val	Pro	Ile	Asn	Pro	Ala	Trp	Arg	Pro
	1				5					10					15	
	Pro	Thr	Val	Thr	Val	Gln	Ala	Gly	Arg	Pro	Ala	Arg	Thr	Pro	Gly	Ala
				20					25					30		
	Pro	Met	Asn	Pro	Pro	Ile	Thr	Leu	Ser	Ser	Thr	Tyr	Val	His	Asp	Ser
20			35					40					45			
	Glu	Lys	Ala	Tyr	Gly	Arg	Asp	Gly	Asn	Asp	Gly	Trp	Gly	Ala	Phe	Glu
	50						55					60				
	Ala	Ala	Met	Gly	Thr	Leu	Asp	Gly	Gly	Phe	Ala	Val	Ser	Tyr	Ser	Ser
	65					70					75					80
25	Gly	Leu	Ala	Ala	Ala	Thr	Ser	Ile	Ala	Asp	Leu	Val	Pro	Thr	Gly	Gly
					85					90					95	
	Thr	Val	Val	Leu	Pro	Lys	Ala	Ala	Tyr	Gly	Val	Thr	Asn	Ile	Phe	
				100					105				110			
	Ala	Arg	Met	Glu	Ala	Arg	Gly	Arg	Leu	Lys	Val	Arg	Thr	Val	Asp	Ala
30			115					120					125			
	Asp	Asn	Thr	Glu	Glu	Val	Ile	Ala	Ala	Ala	Gln	Gly	Ala	Asp	Val	Val
	130						135					140				
	Trp	Val	Glu	Ser	Ile	Ala	Asn	Pro	Thr	Met	Val	Val	Ala	Asp	Ile	Pro
	145					150					155					160
35	Ala	Ile	Val	Asp	Gly	Val	Arg	Gly	Leu	Gly	Val	Leu	Thr	Val	Val	Asp
					165					170					175	
	Ala	Thr	Phe	Ala	Thr	Pro	Leu	Arg	Gln	Arg	Pro	Leu	Glu	Leu	Gly	Ala
				180					185					190		
	Asp	Ile	Val	Leu	Tyr	Ser	Ala	Thr	Lys	Leu	Ile	Gly	Gly	His	Ser	Asp
40			195					200					205			
	Leu	Leu	Leu	Gly	Val	Ala	Val	Cys	Lys	Ser	Glu	His	His	Ala	Gln	Phe
	210															

```

Ala Ile Glu Arg Arg Thr Arg Arg Asp Ala Glu Val Val Ala Gly Val
      340      345      350
Pro Met Thr Leu Cys Arg Val Ser Val Gly Ile Glu Asp Val Glu Asp
      355      360      365
-5 Leu Trp Glu Asp Leu Asn Ala Ser Ile Asp Lys Val Leu Gly
      370      375      380

10 <210> 4
   <211> 32
   <212> DNA
   <213> Artificial sequence

15 <220>
   <223> Description of the artificial sequence: Primer orfR 20

   <400> 4
gatctagaat tcaacatcgt ccgataggcc ag 32

20

   <210> 5
   <211> 40
   <212> DNA
25 <213> Artificial sequence

   <220>
   <223> Description of the artificial sequence: Primer
      orfRmetZ del
30

   <400> 5
ggctctgggta ctggctacga atgcagacaa taatgtcgct 40

35

   <210> 6
   <211> 32
   <212> DNA
   <213> Artificial sequence

40 <220>
   <223> Description of the artificial sequence: Primer metZ 21

   <400> 6
gatctagaat tccgcttccg agaatcaata cc 32

45

   <210> 7
   <211> 402
   <212> DNA
50 <213> Artificial sequence

   <220>
   <223> Description of the artificial sequence: metRmetZ del
      Fragment 1

55

   <400> 7
gatctagaat tcaacatcgt ccgataggcc agcgtctaata tcagcagcga tttcggcagc 60
aagcccagcg ccattaatca gagcgggtgaa ataaacatgg ttcattgatta tgtcaggacg 120
gtaattagac ttatgaccag gtttaaggag gtcaccagggt tgaagccgcg ctattgttcc 180
60 gtggaaaagg gggccctgat ctagctgatt attcatcgca gtaagcgctt tcggttaggtg 240
ggtgaatcat cgtagtcttc cgagccccgt gaccgatcc gttttgtgca atccaatgct 300

```

actcccacag agcgggctac tttctctaaa aatgttctca tagtagataa aattgttctt 360  
aaagcgacat tattgtctgc attcgtagcc agtaaccaga cc 402

5 <210> 8  
<211> 982  
<212> DNA  
<213> Artificial sequence

10 <220>  
<223> Description of the artificial sequence: metRmetZ del  
Fragment 2

<400> 8  
15 gatctagaat tcaacatcgt ccgataggcc agcgtctaatt tcagcagcga tttcggcagc 60  
aagcccagcg ccattaatca gagcgggtgaa ataaacatgg ttcattgatta tgtcaggacg 120  
gtaattagac ttatgaccag gtttaaggag gtcaccaggt tgaagccgag ctattgttcc 180  
gtggaaaagg gggccctgat ctagctgatt attcatcgca gtaagcgctt tcggtaggtg 240  
ggtgaatcat cgtagtcttc cgagccccgt gacccgatcc gttttgtgca atccaatgct 300  
20 actcccacag agcgggctac tttctctaaa aatgttctca tagtagataa aattgttctt 360  
aaagcgacat tattgtctgc attcgtagcc agtaaccaga ctttcagtgt ttggttgcca 420  
ctccagtgtt ggggcccagc gatcagcgaa gttcttcagg atcgacgagt tgatctcaac 480  
acccatttgg ttgggggagg tgagcatcaa ggtgtcggct tccatcacag ctttgtcttc 540  
cttgagctgg tcgatgagtt catcgggact tccggcgtag ctgcgaccga acgtggatcg 600  
25 ggtatcatcc aggattccta cttggtcacc gccttgtccc tgaagtccga aaagctcacg 660  
gtcgcgggtc gtgacgatcg ggaagatgga cctggagaca gacacacgtg ggggtccaatc 720  
gtgtccggct tctttccaag cttggcggtg gaacgcgatt tgatcggtt gcagatcccc 780  
gaaggattgg ccggtggctt cggcgacgag ggtggagctc atcaagttaa cgccatcgcg 840  
tgctgcttgt tcggcggtgt tgtgggaacc ggcgccccac caaatggatt tacgcaggtc 900  
30 aggatcatgc gggaagatcg gcaggggagt gcccggatgg tagaggcgtg ggtattgatt 960  
ctcgggaagcg gaattctaga tc 982

40